



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification n <sup>6</sup> : <b>A61K 31/495, 31/50, 31/44, 31/445</b>	<b>A1</b>	(11) International Publication Number: <b>WO 96/40140</b> (43) International Publication Date: 19 December 1996 (19.12.96)
21) International Application Number: PCT/US96/09561 (22) International Filing Date: 5 June 1996 (05.06.96) (30) Priority Data: 08/474,072           7 June 1995 (07.06.95)           US 08/653,905           28 May 1996 (28.05.96)           US (71) Applicants: GUILFORD PHARMACEUTICALS INC. [US/US]; 6611 Tributary Street, Baltimore, MD 21224 (US). JOHNS HOPKINS UNIVERSITY SCHOOL OF MEDICINE [US/US]; Suite 72-100, 2024 East Monument, Baltimore, MD 21205 (US). (72) Inventors: STEINER, Joseph, P.; 988 Sugar Maple Street, Hampstead, MD 21074 (US). SNYDER, Solomon; Johns Hopkins Medical School, 725 N. Wolfe Street, Baltimore, MD 21205-2185 (US). HAMILTON, Gregory, S.; 6501 Frederick Road, Catonsville, MD 21228 (US). (74) Agent: NATH, Gary, M.; Nath & Associates, 1835 K Street, N.W. #750, Washington, DC 20006-1203 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  Published <i>With international search report.</i> <i>With amended claims.</i>
(54) Title: INHIBITORS OF ROTAMASE ENZYME ACTIVITY		
(57) Abstract  This invention relates to the method of using neurotrophic pipecolic acid derivative compounds having an affinity for FKBP-type immunophilins as inhibitors of the enzyme activity associated with immunophilin proteins, and particularly inhibitors of peptidyl-prolyl isomerase or rotamase enzyme activity to stimulate or promote neuronal growth or regeneration.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

## INHIBITORS OF ROTAMASE ENZYME ACTIVITY

### Related Application

This application is a continuation-in-part application of U.S. Patent Application Serial No. 08/474,072 filed June 7, 1995.

### BACKGROUND OF THE INVENTION

#### 1. Field of the Invention

This invention relates to the method of using neurotrophic pipecolic acid derivative compounds having an affinity for FKBP-type immunophilins as inhibitors of the enzyme activity associated with immunophilin proteins, and particularly inhibitors of peptidyl-prolyl isomerase or rotamase enzyme activity.

#### 2. Description of the Prior Art

The term immunophilin refers to a number of proteins that serve as receptors for the principal immunosuppressant drugs, cyclosporin A (CsA), FK506, and rapamycin. Known classes of immunophilins are cyclophilins, and FK506 binding proteins, such as FKBP. Cyclosporin A binds to cyclophilin while FK506 and rapamycin bind to FKBP. These immunophilin-drug complexes interface with a variety of intracellular signal transduction systems, especially in the immune system and the nervous system.

Immunophilins are known to have peptidyl-prolyl isomerase (PPIase) or rotamase enzyme activity. It has been determined that rotamase activity has a role in the catalyzation of the interconversion of the cis and trans isomer of immunophilin proteins.

Immunophilins were originally discovered and studied in immune tissue. It was initially

postulated by those skilled in the art that inhibition of the immunophilins rotamase activity leads to the inhibition of T-cell proliferation, thereby causing the immunosuppressive action exhibited by immunosuppressive drugs such as cyclosporin A, FK506, and rapamycin. Further study has shown that the inhibition of rotamase activity, in and of itself, is not sufficient for immunosuppressant activity. Schreiber et al. Science 1990, 250, 556-559. Instead immunosuppression appears to stem from the formulation of a complex of immunosuppressant drugs and immunophilins. It has been shown that the immunophilin-drug complexes interact with ternary protein targets as their mode of action. Schreiber et al., Cell 1991, 66, 807-815. In the case of FKBP-FK506 and FKBP-CsA, the drug-immunophilin complexes bind to the enzyme calcineurin, inhibiting T-cell receptor signalling leading to T-cell proliferation. Similarly, the complex of rapamycin and FKBP interacts with the RAFT1/FRAP protein and inhibits signalling from the IL-2 receptor.

Immunophilins have been found to be present at high concentrations in the central nervous system. Immunophilins are enriched 10-50 times more in the central nervous system than in the immune system. Within neural tissues, immunophilins appear to influence nitric oxide synthesis, neurotransmitter release, and neuronal process extension.

Nitric oxide serves several roles in the body. In the brain, nitric oxide appears to be a neurotransmitter. It is formed, from arginine, by nitric oxide synthetase which oxidizes the guanidino group of arginine forming nitric oxide and citrulline. Stimulation of the N-methyl-d-aspartate (NMDA) subtype of glutamate receptors rapidly and markedly activates nitric oxide synthetase and

stimulates cGMP formation. Inhibition of nitric oxide synthetase with arginine derivatives such as nitroarginine blocks the glutamate induced increase in cGMP levels. Nitric oxide synthetase is a  
5 calcium-calmodulin requiring enzyme and N-methyl-d-aspartate receptor activation stimulates nitric oxide synthetase activity because the N-methyl-d-aspartate receptor possesses a calcium channel which is opened by glutamate stimulation, allowing calcium  
10 to rush into the cells and activate the nitric oxide synthetase.

Glutamate is a physiologic neurotransmitter. However, when released in excess, glutamate elicits neurotoxicity via N-methyl-d-aspartate receptors.  
15 Treatment of cerebral cortical neuronal cultures with glutamate or N-methyl-d-aspartate kills up to 90% of neurons and these effects are blocked by N-methyl-d-aspartate antagonist drugs. This N-methyl-d-aspartate neurotoxicity is thought to be a major  
20 contributor to neuronal damage following vascular stroke. Thus, there is a massive release of glutamate following cerebral vascular occlusion and numerous N-methyl-d-aspartate antagonists block stroke damage. Phosphorylation of nitric oxide  
25 synthetase inhibits its catalytic activity. By enhancing nitric oxide synthetase phosphorylation, FK506 might functionally inhibit nitric oxide formation and thus block glutamate neurotoxicity. Indeed, low concentrations of FK506 and cyclosporin  
30 A both block N-methyl-d-aspartate neurotoxicity in cortical cultures. The mediating role of FKBP is evident, as rapamycin reverses the therapeutic effect of FK506. Presumably FK506, already marketed as an immunosuppressant, could be clinically  
35 employed in stroke patients.

FK506 also augments the phosphorylation of growth-associated protein-43 (GAP43). GAP43 is

involved in neuronal process extension and its phosphorylation appears to augment this activity. Accordingly, the effects of FK506 rapamycin and cyclosporin in neuronal process extension have been examined using PC12 cells. PC12 cells are a continuous line of neuronal-like cells which extend neurites when stimulated by nerve growth factor (NGF).

Surprisingly, it has been found that picomolar concentrations of an immunosuppressant such as FK506 and rapamycin stimulate neurite out growth in PC12 cells and sensory nervous, namely dorsal root ganglion cells (DRGs). Lyons et al. *proc. Natl. Acad. Sci. USA*, 1994, 91, 3191-3195. In whole animal experiments, FK506 has been shown to stimulate nerve regeneration following facial nerve injury and results in functional recovery in animals with sciatic nerve lesions.

More particularly, it has been found that drugs with a high affinity for FKBP are potent rotamase inhibitors and exhibit excellent neurotrophic effects. Snyder et al., "Immunophilins and the Nervous System", *Nature Medicine*, Volume 1, No. 1, January 1995, 32-37. These findings suggest the use of immunosuppressants in treating various peripheral neuropathies and enhancing neuronal regrowth in the central nervous system (CNS). Studies have demonstrated that neurodegenerative disorders such as Alzheimer's disease, Parkinson's disease, and amyotrophic lateral sclerosis (ALS) may occur due to the loss, or decreased availability, of a neurotrophic substance specific for a particular population of neurons affected in the disorder.

Several neurotrophic factors effecting specific neuronal populations in the central nervous system have been identified. For example, it has been hypothesized that Alzheimer's disease results from a

decrease or loss of nerve growth factor (NGF). It has thus been proposed to treat Alzheimer's patients with exogenous nerve growth factor or other neurotrophic proteins such as brain derived growth factor, glial derived growth factor, ciliary neurotrophic factor, and neurotrophin-3 to increase the survival of degenerating neuronal populations.

Clinical application of these proteins in various neurological disease states is hampered by difficulties in the delivery and bioavailability of large proteins to nervous system targets. By contrast, immunosuppressant drugs with neurotrophic activity are relatively small and display excellent bioavailability and specificity. However, when administered chronically, immunosuppressants exhibit a number of potentially serious side effects including nephrotoxicity, such as impairment of glomerular filtration and irreversible interstitial fibrosis (Kopp et al., 1991, *J. Am. Soc. Nephrol.* 1:162); neurological deficits, such as involuntary tremors, or non-specific cerebral angina such as non-localized headaches (De Groen et al., 1987, *N. Engl. J. Med.* 317:861); and vascular hypertension with complications resulting therefrom (Kahan et al., 1989 *N. Engl. J. Med.* 321: 1725).

The present invention provides non-immunosuppressive as well as immunosuppressive pipecolic acid derivative compounds containing small molecule FKBP rotamase inhibitors which are extremely potent in augmenting neurite outgrowth, and for promoting neuronal growth, and regeneration in various neuropathological situations where neuronal repair can be facilitated including peripheral nerve damage by physical injury or disease state such as diabetes, physical damage to the central nervous system (spinal cord and brain), brain damage associated with stroke, and for the

treatment of neurological disorders relating to neurodegeneration, including Parkinson's disease, Alzheimer's disease, and amyotrophic lateral sclerosis.

5

#### SUMMARY OF THE INVENTION

This invention relates to the method of using neurotrophic pipecolic acid derivative compounds having an affinity for FKBP-type immunophilins as inhibitors of the enzyme activity associated with immunophilin proteins, and particularly inhibitors of peptidyl-prolyl isomerase or rotamase enzyme activity.

A preferred embodiment of this invention is a method of treating a neurological disorder in an animal, comprising:

administering to an animal an effective amount of a pipecolic acid derivative having an affinity for FKBP-type immunophilins to stimulate growth of damaged peripheral nerves or to promote neuronal regeneration, wherein the FKBP-type immunophilin exhibits rotamase activity and the pipecolic acid derivative inhibits said rotamase activity of the immunophilin.

Another preferred embodiment of this invention is a method of treating a neurological disorder in an animal, comprising:

administering to an animal an effective amount of a pipecolic acid derivative having an affinity for FKBP-type immunophilins in combination with an effective amount of a neurotrophic factor selected from the group consisting of neurotrophic growth factor, brain derived growth factor, glial derived growth factor, ciliary neurotrophic factor, and neurotrophin-3, to stimulate growth of damaged



peripheral nerves or to promote neuronal regeneration, wherein the FKBP-type immunophilin exhibits rotamase activity and the pipecolic acid derivative inhibits said  
5 rotamase activity of the immunophilin.

Another preferred embodiment of this invention is a method of stimulating growth of damaged peripheral nerves, comprising;

10 administering to damaged peripheral nerves an effective amount of a pipecolic acid derivative compound having an affinity for FKBP-type immunophilins to stimulate or promote growth of the damaged peripheral nerves, wherein the FKBP-type immunophilins exhibit rotamase  
15 activity and the pipecolic acid derivative inhibits said rotamase activity of the immunophilin.

Another preferred embodiment of this invention is a method of stimulating growth of damaged  
20 peripheral nerves, comprising:

administering to damaged peripheral nerves an effective amount of a pipecolic acid derivative compound having an affinity for FKBP-type immunophilins to stimulate growth of damaged  
25 peripheral nerves, wherein the FKBP-type immunophilin exhibit rotamase activity and the pipecolic acid derivative inhibits said rotamase activity of the immunophilin.

Another preferred embodiment of this invention is a method for promoting neuronal regeneration and  
30 growth in animals, comprising:

administering to an animal an effective amount of a pipecolic acid derivative compound having an affinity for FKBP-type  
35 immunophilins to promote neuronal regeneration, wherein the FKBP-type immunophilins exhibit rotamase activity and the pipecolic acid

derivative inhibits said rotamase activity of the immunophilin.

Yet another preferred embodiment of this invention is a method for preventing

5 neurodegeneration in an animal, comprising:  
administering to an animal an effective amount  
of a pipecolic acid derivative having an  
affinity for FKBP-type immunophilins to prevent  
neurodegeneration, wherein the FKBP-type  
10 immunophilin exhibits rotamase activity and the  
pipecolic acid derivative inhibits said  
rotamase activity of the immunophilin.

#### BRIEF DESCRIPTION OF THE DRAWINGS

##### Figure 1

15 FKBP-12 and GAP-43 expression in the facial  
nucleus after nerve crush. *In situ* hybridization  
comparing the time course of expression of mRNA in  
the facial nucleus for FKBP12 (left) and GAP-43  
(right). The right facial nucleus is ipsilateral to  
20 the crush, and the left side is an unoperated  
control (Fig 1B). *In situ* hybridization for FKBP-12  
on an untreated control (left) and for calcineurin  
 $A\alpha, \beta$  7 days following facial nerve crush (right).

25 Experiments were replicated at least 3 times  
with similar results.

##### Figure 2

Localization of FKBP-12 to facial motor neurons  
following nerve crush. Bright-field  
photomicrographs of *in situ* hybridization for FKBP-  
30 12 in motor neurons of the facial nucleus 7 days  
after crush (Fig. 2A), and in motor neurons of  
control facial nucleus (Fig. 2B).

##### Figure 3

35 Upregulation of FKBP-12 mRNA in lumbar spinal  
cord motor neurons after sciatic nerve crush. *In situ*  
hybridization for FKBP-12 7 days after crush

of the right sciatic nerve. Top panel (Fig. 3A) shows the response of motor neurons in the ventral horn of lower lumbar spinal cord (indicated by the arrow). Bright field photomicrographs of  
5 corresponding motor neuron pools are shown in the bottom panels: (Fig. 3B) left side contralateral to nerve crush, (Fig. 3C) right side ipsilateral to the nerve crush. This experiment was repeated 3 times with similar results.

10           **Figure 4**

Induction of FKBP and FKBP-12 mRNA in the dorsal root ganglion 1 and 6 weeks after sciatic nerve crush. Dark-field photomicrographs of  
15 sections through the L4 dorsal root ganglion ipsilateral to sciatic nerve crush processed for FKBP in situ hybridization are shown in the left panels and for [<sup>3</sup>H]FK506 autoradiography in the right panels. These results were replicated 3 times for each time point.

20           **Figure 5**

Ricin lesion of the right facial nerve. Nissl stain (bottom panel, Fig. 5A) reveals extensive degeneration of motor neurons in the right facial nucleus with an accompanying glial proliferation 7  
25 days following injection of ricin into the facial nerve. In situ hybridization for FKBP mRNA 7 days after ricin lesion of the facial nerve/nucleus is shown in the top panel (Fig. 5B). This experiment was replicated 3 times with similar results.

30           **Figure 6**

[<sup>3</sup>H]FK506 binding in segments of sciatic nerve 7 days following crush. The diagram illustrates the 3  
35 mm segments of nerve taken: constrictions indicate positions of ligatures applied at day 7 for the 6 hr collection time as described in the methods. The distal ligature site is 10 mm proximal to the original crush site. Anterograde transport of FKBP

is 124 mm/day. Data are the means  $\pm$  S.E.M. (n = 3).

#### Figure 7

Transport of FKBP in the sciatic nerve. Dark-field photomicrographs of sections through a control (untreated) sciatic nerve and a 7 day sciatic nerve crush site processed for FKBP-12 *in situ* hybridization (Fig. 7A, Fig. 7B) and for [<sup>3</sup>H]FK-506 autoradiography (Fig. 7C, Fig. 7D). Arrows indicate the sight of the nerve crush. This experiment was repeated 3 times with similar results.

#### Figure 8

Levels of [<sup>3</sup>H]FK506 binding in PC-12 cells maintained in the presence or absence of NGF (50 ng/ml). n = 3 for each time point. Bars represent S.E.M.

#### Figure 9

Immunosuppressant mediated enhancement of neurite outgrowth in PC-12 cells. Hoffman contrast photomicrographs (64) of cultures grown for 48 hr in the presence of NGF with or without added FK506 or rapamycin. Fig. 9A: PC-12 cells grown in 1.0 ng/ml NGF. Fig. 9B: 50 ng/ml NGF. Fig. 9C: 1.0 ng/ml NGF and 100 nM FK506. Fig. 9D: 1.0 ng/ml NGF and 100 nM rapamycin. Magnification 200 X.

#### Figure 10

Effects of FK506 on neurite outgrowth in PC-12 cells. Cultures were treated with varying concentrations of NGF in the presence or absence of 100 nM FK506, and neurite sprouting was measured at 48 hr. Outgrowth was quantitated as described in Methods by counting cells with neuritic processes greater than 5  $\mu$ m. n = 4 separate experiments for each point and error bars represent SEM.

#### Figure 11

Concentration-response relationship for FK506 potentiation of neurite outgrowth in PC-12 cells. Cells were treated for 48 hr with 1 ng/ml NGF and

varying concentrations of FK506. Neurite outgrowth response was measured as described in Figure 10 and Methods. n = 4 separate experiments for each data point \*p<.001 Students t test.

5                   Figure 12

[<sup>3</sup>H]FK-506 autoradiography on dorsal root ganglion explant cultures. After 26 days of cultures with 100 ng/ml NGF the extensive processes display abundant FKBP associated silver grains. 10       Autoradiographic grains are abolished with 1  $\mu$ M unlabeled FK506.

                  Figure 13

Phase-contrast micrographs of dorsal root ganglia grown with different substances. Fig. 13A: 15       NGF 100 ng/ml, Fig. 13B: FK506 1  $\mu$ M, Fig. 13C: FK506 1  $\mu$ M and anti-NGF antibody, Fig. 13D: No added growth factor, Fig. 13E: FK506 1pM, Fig. 13F: FK506 1 $\mu$ M. and rapamycin 1  $\mu$ M. Scale bar is 205  $\mu$ m. NGF produces abundant axon outgrowth (Fig. 13A), as does 20       1  $\mu$ M FK506 (Fig. 13B). The effects of FK506 are substantially decreased by reducing the concentration to 1 pM (Fig. 13E). However, neurite outgrowth with 1 pM FK506 is greater than in its absence (Fig. 13D). FK506 effects are also 25       diminished by adding anti-NGF antibody to eliminate the effects of NGF produced by non-neuronal cells in the cultures. The abundant neurites that occur in large fascicles in response to NGF (100 ng/ml)] (Fig. 13A) or 1  $\mu$ M FK506 (Fig. 13B) appear white, 30       while small fascicles or individual neurites appear black. Non-neuronal cells, Schwann cells and some fibroblasts, are more evident with 1 pM FK506 (Fig. 13E) or anti-NGF antibody (Fig. 13C) than with 1  $\mu$ M FK506 (Fig. 13B). NGF produced by non-neuronal 35       cells in the cultures results in the limited axon outgrowth seen in cultures with no added growth factors (Fig. 13D). The large number of refractile

non-neuronal cells, appearing white, tend to overshadow the few neurites (Fig. 13D). Rapamycin completely inhibits axon outgrowth in the presence of FK506 (Fig. 13F). Micrographs are representative of 12-30 ganglia from each experimental condition. Differences between all experimental groups were highly reproducible.

#### Figure 14

Effects of FK506 and rapamycin on NGF-mediated neurite extension in PC12 cells. PC12 cells (passage 60) were treated with various concentrations of NGF alone or in the presence of 100 nM FK506, 100 nM rapamycin or 100 nM WAY-124,466. Neurite outgrowth was measured after 96 hours with cells bearing processes longer than the diameter of the cells scoring positive. n=3 separate experiments for each point and error bars represent S.E.M.

#### Figure 15

Picomolar concentrations of (A) FK506 and (B) rapamycin and WAY-124,466 potentiate neurite extension elicited by NGF (0.5 ng/ml) in PC12 cells. Low passage PC12 cells were treated for 4 days with 0.5 ng/ml NGF in the presence of various concentrations of FK506 ( $\square$ ), rapamycin ( ) or WAY-124,466 ( ). Neurite expression was quantitated as described above in figure 14. The levels of neurite production in the presence of 0.5 ng/ml NGF (designated L) and 50 ng/ml NGF (designated H) are indicated for comparative purposes.

#### Figure 16

Photomicrographs of PC12 cells treated with immunophilin ligands + 0.5 ng/ml NGF itself or 50 ng/ml NGF.

#### Figure 17

Immunophilin ligands reduce the amount of NGF

required to produce maximal neurite extension in chick sensory ganglia. Whole dorsal root ganglion explants were isolated from day 9-10 chick embryos and cultured in Matrigel-coated 12-well dishes containing L15 medium plus high glucose, with 10% fetal calf serum supplemented with 10  $\mu$ M Ara C (penicillin and streptomycin) at 37°C in a 5% CO<sub>2</sub> environment. Sensory ganglia were treated with 1 ng/ml NGF, 1 ng/ml NGF plus 100 nM FK506 or 100 ng/ml NGF for 48 hr, and neuronal processes were counted and photographed.

#### Figure 18

FK506, rapamycin, and WAY-124,466 potentiate NGF-dependent neurite production in sensory ganglia. Explants of chick DRG were cultured as described in figure 17 above. FK506, rapamycin and WAY-124,466 (100nM each plus or minus 0.1 ng/ml NGF were added to the DRG explant cultures. At 48 hrs., neurite outgrowth was quantitated and the cultures were photographed.

#### Figure 19

Photomicrograph of Example 111 promoting neurite outgrowth in Chick dorsal root ganglion cultures. The three panels show neurite outgrowth at 1 pM concentration (left panel), 100 pM concentration (center panel), and 100 nM concentration (right panel) of Example 111.

#### Figure 20

Photomicrograph of Example 17 promoting neurite outgrowth in dorsal root ganglion cultures. The three panels show neurite outgrowth at 1 pM concentration (left panel), 100 pM concentration (center panel), and 100 nM concentration (right panel) of Example 17.

Figure 21

Photomicrograph of Example 102 promoting neurite outgrowth in dorsal root ganglion cultures. The three panels show neurite outgrowth at 1 pM concentration (left panel), 100 pM concentration (center panel), and 100 nM concentration (right panel) of Example 102.

#### DETAILED DESCRIPTION OF THE INVENTION

The novel neurotrophic pipecolic acid derivative compounds of this invention have an affinity for the FK506 binding proteins such as FKBP-12. When the neurotrophic compounds of the invention are bound to FKBP, they have been found to inhibit the prolyl- peptidyl cis-trans isomerase activity, or rotamase activity of the binding protein and unexpectedly stimulate neurite growth.

The compounds of the present invention can be used in the form of salts derived from inorganic or organic acids and bases. Included among such acid salts are the following: acetate, adipate, alginate, aspartate, benzoate, benzenesulfonate, bisulfate butyrate, citrate, camphorate, camphorsulfonate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, fumarate, glucoheptanoate, glycerophosphate, hemissulfate heptanoate, hexanoate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethanesulfonate, lactate, maleate, methanesulfonate, 2-naphthalensulfonate, nicotinate, oxalate, pamoate, pectinate, propionate, succinate, tartrate, thiocyanate, tosylate and undecanoate. Base salts include ammonium salts, alkali metal salts such as sodium and potassium salts, alkaline earth metal salts such as calcium and magnesium salts, salt with organic bases such as dicyclohexylamine salts, N-



methyle-D-glucamine, and salts with amino acids such as arginine, lysine, and so forth. Also, the basic nitrogen-containing groups can be quarternized with such agents as lower alkyl halides, such as methyl, ethyl, propyl, and butyl chloride, bromides and iodides; dialkyl sulfates like dimethyl, diethyl, dibutyl and diamyl sulfates, long chain halides such as decyl, lauryl, myristyl and stearyl chlorides, bromides and iodides, aralkyl halides like benzyl and phenethyl bromides and others. Water or oil-soluble or dispersible products are thereby obtained.

The neurotrophic compounds of this invention can be periodically administered to a patient undergoing treatment for neurological disorders or for other reasons in which it is desirable to stimulate neuronal regeneration and growth, such as in various peripheral neuropathic and neurological disorders relating to neurodegeneration. The compounds of this invention can also be administered to mammals other than humans for treatment of various mammalian neurological disorders.

The novel compounds of the present invention are potent inhibitors of rotamase activity and possess an excellent degree of neurotrophic activity. This activity is useful in the stimulation of damaged neurons, the promotion of neuronal regeneration, the prevention of neurodegeneration, and in the treatment of several neurological disorders known to be associated with neuronal degeneration and peripheral neuropathies. The neurological disorders that may be treated include but are not limited to: trigeminal neuralgia, glossopharyngeal neuralgia, Bell's Palsy, myasthenia gravis, muscular dystrophy, amyotrophic lateral sclerosis, progressive muscular atrophy, progressive bulbar inherited muscular atrophy,

herniated, ruptured or prolapsed intervertebral disk syndromes, cervical spondylosis, plexus disorders, thoracic outlet destruction syndromes, peripheral neuropathic such as those caused by lead, dapsone, ticks, porphyria, or Gullain-Barré syndrome, Alzheimer's disease, and Parkinson's disease.

For these purposes the compounds of the present invention may be administered orally, parenterally, by inhalation spray, topically, rectally, nasally, buccally, vaginally or via an implanted reservoir in dosage formulations containing conventional non-toxic pharmaceutically-acceptable carriers, adjuvants and vehicles. The term parenteral as used herein includes subcutaneous, intravenous, intramuscular, intraperitoneally, intrathecally, intraventricularly, intrasternal and intracranial injection or infusion techniques.

To be effective therapeutically as central nervous system targets the immunophilin-drug complex should readily penetrate the blood-brain barrier when peripherally administered. Compounds of this invention which cannot penetrate the blood-brain barrier can be effectively administered by an intraventricular route.

The pharmaceutical compositions may be in the form of a sterile injectable preparation, for example as a sterile injectable aqueous or oleaginous suspension. This suspension may be formulated according to techniques known in the art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium

chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. Fatty acids such as oleic acid and its glyceride derivatives find use in the preparation of injectables, olive oil or castor oil, especially in their polyoxyethylated versions. These oil solutions or suspensions may also contain a long-chain alcohol diluent or dispersant.

The compounds may be administered orally in the form of capsules or tablets, for example, or as an aqueous suspension or solution. In the case of tablets for oral use, carriers which are commonly used include lactose and corn starch. Lubricating agents, such as magnesium stearate, are also typically added. For oral administration in a capsule form, useful diluents include lactose and dried corn starch. When aqueous suspensions are required for oral use, the active ingredient is combined with emulsifying and suspending agents. If desired, certain sweetening and/or flavoring and/or coloring agents may be added.

The compounds of this invention may also be administered in the form of suppositories for rectal administration of the drug. These compositions can be prepared by mixing the drug with a suitable non-irritating excipient which is solid at room temperature but liquid at rectal temperature and therefore will melt in the rectum to release the drug. Such materials include cocoa butter, beeswax and polyethylene glycols.

The compounds of this invention may also be administered optically, especially when the conditions addressed for treatment involve areas or organs readily accessible by topical application, including neurological disorders of the eye, the

skin, or the lower intestinal tract. Suitable topical formulations are readily prepared for each of these areas.

For ophthalmic use, the compounds can be formulated as micronized suspensions in isotonic, pH adjusted sterile saline, or, preferably, as solutions is isotonic, pH adjusted sterile saline, either with or without a preservative such as benzylalkonium chloride. Alternatively for the ophthalmic uses the compounds may be formulated in an ointment such as petrolatum.

For application topically to the skin, the compounds can be formulated in a suitable ointment containing the compound suspended or dissolved in, for example, a mixture with one or more of the following: mineral oil, liquid petrolatum, white petrolatum, propylene glycol, polyoxyethylene polyoxypropylene compound, emulsifying wax and water. Alternatively, the compounds can be formulated in a suitable lotion or cream containing the active compound suspended or dissolved in, for example, a mixture of one or more of the following: mineral oil, sorbitan monostearate, polysorbate 60, cetyl esters wax, cetearyl alcohol, 2-octyldodecanol, benzyl alcohol and water.

Topical application for the lower intestinal tract an be effected in a rectal suppository formulation (see above) or in a suitable enema formulation.

Dosage levels on the order of about 0.1 mg to about 10,000 mg. of the active ingredient compound are useful in the treatment of the above conditions, with preferred levels of about 0.1 mg. to about 1,000 mg. The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration.

It is understood, however, that a specific dose level for any particular patient will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, rate of excretion, drug combination, and the severity of the particular disease being treated and form of administration.

The compounds can be administered with other neurotrophic agents such as neurotrophic growth factor (NGF), glial derived growth factor, brain derived growth factor, ciliary neurotrophic factor, and neurotrophin-3. The dosage level of other neurotrophic drugs will depend upon the factors previously stated and the neurotrophic effectiveness of the drug combination.

#### Methods and Procedures

##### Sciatic Nerve Crush

This example demonstrates high levels of FKBP in normal peripheral nerve and that these increase following nerve crush.

If FKBP were physiologically associated with neuronal process extension in the actions of GAP-43, then one might anticipate substantial levels of FKBP in peripheral nerve. Accordingly, we measured [<sup>3</sup>H]FK-506 binding in rat sciatic nerve, as well as in growth cones isolated from 2-day-old rat pups, and compared values with those of the cerebral cortex and several peripheral tissues.

[<sup>3</sup>H]FK-506 autoradiography was carried out as described on unfixed sections which were thawed and air dried before preincubation for 1 hr in buffer consisting of 50 mM Hepes, 2 mg/ml bovine serum albumin, 5% ethanol, and 0.02% Tween 20 pH 7.4. Sections were then exposed to 1 nM [<sup>3</sup>H]FK-506 (86.5 Ci/mMol; DuPont-NEN, Boston, MA) for 1 hr at room temperature in preincubation buffer. Non-specific

binding was defined by addition of 1  $\mu$ M FK-506. Following incubation, the slides were washed 4 x 5 min in ice cold preincubation buffer and air dried. The radiolabeled sections were then juxtaposed to tritium-sensitive film or coverslips coated with Kodak NTB-2 emulsion.

TABLE 1

[<sup>3</sup>H]FK506 Binding to Sciatic Nerve and Growth Cones

(A) [<sup>3</sup>H]FK506 Binding in Sciatic Nerve

Tissue	Bmax (pmol/mg protein)
Adult Rat	
Sciatic Nerve	22.1
Cerebral Cortex	38.0
Thymus	9.5
Spleen	8.0
Neonatal Rat	
Forebrain	25.5
Growth Cones	10.2

(B) [<sup>3</sup>H]FK506 Binding After Sciatic Nerve Crush

protein	Bmax fmol/5mm segment	Bmax pmol/mg
Unoperated	31.8 $\pm$ 2.1	21.2 $\pm$ 1.4
7-Day Crush	136.5 $\pm$ 15.7*	40.1 $\pm$ 2.0*

[<sup>3</sup>H]FK506 binding was assayed as described in methods. In Table 1A experiments were replicated three times with less than 10% variation. In Table 1B values are presented as the mean  $\pm$  S.E.M. (n = 3). \*P<sub>≤</sub> 0.05 Students' t-test for independent means.

Of all the tissues examined sciatic nerve binding levels are the highest, somewhat higher than those of the cerebral cortex and about 10 x higher

than levels in the thymus and spleen, which contain FKBP associated with lymphocytes. See Table 1A.

Evidence for a role of FKBP in nerve regeneration comes from experiments in which we  
5 crushed the sciatic nerve of adult rats and 7 days later measured [<sup>3</sup>H]FK506 binding in a 5 mm segment immediately proximal to the nerve crush.

Sprague-Dawley rats (175-200g) were anesthetized with a mixture of Rompun (12 mk/kg)  
10 Ketamine (30 mg/kg). Using aseptic techniques, the facial nerve was crushed with jewelers forceps 2 x 30 sec 2 mm distal to its exit from the stylomastoid foramen. Identical procedures were used to crush the sciatic nerve at the level of the mid-thigh.

15 Total binding in the segment proximal to the crush was quadrupled compared to control values. Since total protein is substantially augmented in the proximal segment, [<sup>3</sup>H]FK-506 binding per mg protein is only doubled in the proximal segment.

#### 20 Facial Nerve Crush

This example demonstrates that facial nerve lesions augment the coincident expression of FKBP and GAP-43.

Following the crush of the facial nerve, mRNA  
25 levels of GAP-43 increase in the facial nerve nucleus. Utilizing *in situ* hybridization, we examined mRNA levels of FKBP, GAP-43 and calcineurin following facial nerve crush.

Rats were perfused transcardially with 150-200 ml ice cold phosphate-buffered saline (PBS) (0.1M, pH 7.4). Tissues were removed and immediately frozen in isopentane (-80° C). Cryostat sections 18  
5  $\mu$ m thick) were cut and thaw mounted on gelatin coated slides.

In situ hybridization was performed as previously described, using antisense oligonucleotide probes end labeled with [<sup>35</sup>S]dATP.  
10 For FKBP, three separate oligonucleotides complementary to the following regions of the cloned cDNA disclosed by Maki, et al. (1990) *Proc. Natl. Acad.-Sci. USA* 87, 5440-5443, and Standaert, R.F., et al. (1990) *Nature* 346, 671-674 and incorporated  
15 herein by reference, were used: 70-114, 214-258, 441-485. For GAP-43, three separate antisense oligonucleotides complementary to nucleotides 961-1008, 1081-1128, 1201-1248 of the cloned cDNA disclosed by Rosenthal, A., et al. (1987) *EMBO J.* 6,  
20 3641-3646 and incorporated herein by reference were used. For calcineurin A $\alpha$  antisense oligonucleotides complementary to the nucleotides 1363-1410 and 1711-1758, disclosed by Ito et al. (1989) *Biochem. Biophys. Res. Commun.* 163, 1492-1497 and  
25 incorporated herein by reference, and for calcineurin A $\beta$  1339-1386 and 1569-1616 disclosed by Kuno, T., et al. (1989) *Biochem. Biophys. Res. Commun.* 165, 1352-1358 and incorporated herein by



reference, were used. Sections were thawed and allowed to dry, then fixed for 5 min in 4% freshly depolymerized paraformaldehyde in PBS. Following two rinses in PBS, sections were acetylated with  
5 0.25% acetic anhydride in 0.1 M triethanolamine 0.5% NaCl (pH 8.0), and then dehydrated in graded alcohols, defatted in chloroform for 5 min, rehydrated to 95% ethanol and allowed to air dry. Hybridization was performed overnight at 37° C in  
10 buffer containing 50% deionized formamide, 10% dextran sulfate, 4 x SSC, 1 x Denhardt's solution, 20 mM phosphate buffer, 0.1 mg/ml salmon sperm DNA, 0.1 mg/ml yeast transfer RNA, 10 mM dithiothreitol, 2.0% betamercaptoethanol (BMD), 1.0 mM EDTA and  
15 labelled probe (2,000,000 dpm/section). Following hybridization, sections were rinsed in 1 x SSC, 1.0% BME for 15 min at room temperature, then twice for 10 min at 55° C air dried and placed on film or dipped in Kodak NTB-2 emulsion.

20 Striking enhancement of FKBP and GAP-43 expression is observed, while no changes are evident in calcineurin expression. As early as 24 hr following facial nerve crush FKBP expression is increased with peak levels evident a 1-2 weeks,  
25 while mRNA concentrations diminish substantially at 3 weeks. Examination under higher magnification reveals that the increased levels of silver grains for FKBP mRNA are confined to neuronal cell bodies

(Figure 2). Northern blot analysis of the dissected facial nucleus confirms the increased levels of FKBP specific mRNA. GAP-43 mRNA levels follow a time course closely similar to those of FKBP. By contrast, no changes in calcineurin expression are detected at any of the time points examined.

Total cellular RNA from the dissected facial nucleus was isolated. Samples of 10 or 20 ug total RNA were electrophoresed through a 1% agarose, 2.0% formaldehyde gel and transferred to a nylon membrane in 10 nM NaOH. cDNA probes to FKBP labeled with [<sup>35</sup>]dCTP to a specific activity of  $1 \times 10^9$  cpm/ug by random priming were hybridized overnight at 42° C in buffer consisting of 50% formamide, 2 x SSPE, 7% SDS, 0.5% Blotto and 100 ug/ml salmon sperm DNA. The blots were washed for 20 min at room temperature, and 2 x 15 min at 65° C in 0.15 x SSC, 0.15% SDS and then exposed to film for 48096 hrs.

On the unlesioned side a modest increase in silver grains compared to control sections are observed. This is consistent with findings that contralateral neurons also respond to axotomy.

Following facial nerve crush, rats develop a facial nerve palsy, which is evident by the lack of whisker movement with functional recovery at 3 weeks coincident with the completion of nerve regeneration. In our rats we also observed the loss of whisker movement following nerve crush with a

return of function at 3 weeks. Thus, the time course of increased expression of GAP-43 and FKBP correlates with the process of nerve regeneration.

#### Sciatic Nerve Regeneration

5           This example demonstrates alterations in FKBP and GAP-43 associated with sciatic nerve regeneration.

          Following sciatic nerve lesions GAP-43 mRNA levels are enhanced in both spinal cord motor  
10           neurons and in dorsal root ganglia neuronal cells. In rats subjected to sciatic nerve crush, we observed a striking enhancement in mRNA levels for FKBP in motor neurons at L-4, 5 (Figure 3) and in dorsal root ganglia neuronal cells coincident with  
15           the reported enhancement of GAP-43 expression (Figure 4). At high magnification we observed the FKMB mRNS silver grains localized to neuronal cell bodies (Figure 3). We monitored FKBP protein levels by autoradiography of [<sup>3</sup>H]FK-506 binding under  
20           conditions in which it binds selectively to FKBP (Figure 4). Increased FKBP is detected in the primary sensory neurons in the dorsal root ganglia, though no increases are evident in motor neuronal cells following sciatic nerve crush.

25           The association of augmented FKBP expression with regeneration selectively is further supported by experiments with ricin. When injected into peripheral nerves ricin is transported back into the

cell body which is destroyed without associated nerve regeneration. We injected 0.5 ug ricin (R 60, Sigma, St. Louis, MO) into the facial ne : the same site where crushes had been performed in other experiments according to the method of Str and Kreutzberg in 0.5 ul PBS and 0.1% Fast Gree Streit et al., (1988) *J. Comp. Neurol.* 268, 248-

We conducted in situ hybridization localization studies for FKBP mRNA at 2, 4 and 7 days following ricin treatment (Figure 5). No increase in FKBP mRNA is observed following ricin treatment. Gli occurs both following ricin treatment and nerve crush. The failure of FKBP mRNA to increase following ricin treatment fits with the selective neuronal localization of FKBP mRNA in the facial nucleus.

#### FKBP Transport in the Sciatic Nerve

This example demonstrates that FKBP is rapidly transported in the sciatic nerve.

The failure of FKBP protein to increase in motor neurons following sciatic nerve crush despite the increase in FKBP mRNA suggests that the protein is rapidly transported out of the cell body into nerve processes. This fits with our earlier observations that FKBP mRNA is concentrated in granule cells of the cerebellum which contain low levels of FKBP protein, while FKBP protein levels are highly concentrated in the molecular layer.

the cerebellum associated with the parallel fibers arising from granule cells. To examine for possible transport of FKBP, we crushed the sciatic nerve and 7 days later applied ligatures 10 and 20 mm proximal to the crush. Six hr following ligature, we monitored [<sup>3</sup>H]FK-506 binding in 3 mm segments spanning the area of the ligatures. (Figure 6).

For axon transport experiments, classic ligature techniques were used following the methods of Tetzlaff et al. One week following sciatic nerve crush two collection ligatures (510 sutures) were placed on the nerve approximately 10 mm apart with the distal most ligature positioned 10 mm proximal to the initial crush site. Six hours later, 5-3 mm segments of the nerve were collected from regions proximal to, distal to, and between the collection ligatures as illustrated in Figure 5. The nerve segments were prepared for [<sup>3</sup>H]FK-506 binding assays by homogenizing in 10 volumes of 50 mM Tris-HCl, pH 7.4. Homogenates were centrifuged at 15,000 x g for 20 min at 4° C, and supernatants were collected and assayed for total protein concentration using the Coomassie blue dye binding assay (Pearce). [<sup>3</sup>H]FK-506 binding was carried out as described (4) on aliquots containing 2 ug of total soluble protein in a final volume of 0.4 ml assay buffer consisting of 50 mM Tris-HCl, pH 7.4, 2 mg/ml bovine serum albumin, 250 pM [<sup>3</sup>H]FK-506, and varying

concentrations of unlabeled FK-506. Following incubation at 25° C for 60 min, 0.35 ml was layered over a 0.8 ml column of LH-20 Sephadex (Pharmacia LKB) and washed with 0.4 ml of assay buffer. The  
5 eluates were collected and counted in a scintillation counter.

Results are shown in Figure 5. [<sup>3</sup>H]FK-506 binding levels are highest in the segment just proximal to the ligature 20 cm from the crush, being  
10 almost quadruple levels in the other segments. Based on the levels of [<sup>3</sup>H]FK-506 binding in segments A-D, we calculated the rate of anterograde transport for FKBP. This rate of 240 mm per day is  
essentially the same as transport rates for GAP-43  
15 representing the most rapid transport rates for neuronal proteins.

To visualize the accumulation of FKBP following nerve crush, we applied a loose ligature to mark the site of crush of the sciatic nerve and conducted in  
20 situ hybridization for FKBP mRNA as well autoradiography for [<sup>3</sup>H]FK-506 binding (Figure 7). Most FKBP mRNA and [<sup>3</sup>H]FK-506 binding accumulate immediately proximal to the crush. These levels are considerably higher than in control uncrushed  
25 sciatic nerve. Examination of the in situ hybridization an autoradiography preparations at high magnification reveals silver grains associated with neuronal fibers. There are also silver grains

localized to cells whose identity we could not determine definitively, so that they may be Schwann cells, macrophages or fibroblasts.

#### FKBP in PC12 Cells

5           This example demonstrates that PC-12 cells contain FKBP and that FKBP levels are enhanced by nerve growth factor. We examined PC-12 cells for the presence of FKBP by monitoring the binding of [3H]FK-506 to cells under basal conditions and  
10           following treatment with nerve growth factor (NGF).

          Levels of FKBP in PC-12 cells were obtained from Scatchard analysis of [3H]FK-506 binding curves. Cultures were scraped from the culture wells and homogenized in 10 volumes of 50 mM Tris-HCl, pH 7.4,  
15           1 mM EDTA, 100 µg/ml phenylmethylsulfonylfluoride and centrifuged at 40,000 x g for 20 min at 4° C. Protein was determined by the Coomassie blue dye binding assay using bovine serum albumin as a standard. Binding of 250 pM [3H]dihydro FK506 (86.5  
20           Ci/mmol, DuPont/NEN) was assessed for samples containing 5 µg soluble protein in a final volume of 0.4 ml assay buffer containing 50 mM Tris-HCl, pH 7.4, 2 mg/ml BSA and varying concentrations of unlabeled FK506. After 60 min incubation at 25° C,  
25           0.35 ml was layered over a 0.8 ml column of LH-20 Sephadex (Pharmacia LKB), pre-equilibrated with assay buffer. The column was further washed with 0.4 ml of assay buffer, the eluates collected, mixed

with Formula 963 (DuPont/NEN) and counted in a Beckman scintillation counter. Specific binding was determined by subtracting binding obtained in the presence of 1  $\mu$ M unlabeled FK506 from total [ $^3$ H]FK506 bound.

Results are shown in Figure 8. [ $^3$ H]FK506 binds saturably to untreated PC-12 cell homogenates. In typical experiments about 1,000 cpm are bound while nonspecific binding in the presence of 1  $\mu$ M FK506 is about 150 cpm. Fifty percent inhibition of [ $^3$ H]FK506 binding occurs with 1-2 nM FK506 indicating that the binding sites correspond to authentic FKBP.

[ $^3$ H]FK506 binding increases markedly following NFG treatment. Significant increases are evident by 10-15 hr. Binding triples by 20 hr and a modest further increase is evident at 100 hr.

#### Increased Neurite Extension in PC12 Cells

This example demonstrates that FK506 and rapamycin increase neurite extension in PC-12 cells.

PC-12 cells were maintained at 37° C, 5% CO<sub>2</sub>, in Dulbecco's modified Eagle medium (DMEM) supplemented with 10% heat-inactivated horse serum and 5% heat-inactivated fetal bovine serum. For differentiation in NGF, cells were plated at 1 x 10<sup>5</sup> in 35 mm culture wells coated with rat tail collagen at 5  $\mu$ g/cm<sup>2</sup>, and allowed to attach before replacing the media with DMEM supplemented with 2% fetal horse serum, NGF and/or FK506 or rapamycin. For quantitation of



neurite outgrowth, random photographs were made (3-4 per well), and process bearing neurons were counted with processes being greater than 5  $\mu$ m.

Experimental conditions were unknown by the  
5 photographer and cell counter. Four separate experiments were performed in duplicate for each data point presented. Neurites were identified and counted from approximately 100 cells per photograph. Thus, neurites from 1200 - 1600 cells were counted  
10 per data point.

As observed, NGF potently stimulates neurite outgrowth with half-maximal stimulation at 1 ng/ml and maximal augmentation at about 50-100 ng/ml (Figs. 9, 10). FK506 (100 nM) markedly augments the  
15 effect of NGF by increasing sensitivity to NGF. Thus, FK506 reduces by 20-50 fold the NGF concentration needed to elicit maximal outgrowth. Half maximal outgrowth in the absence of FK506 occurs at 5 ng/ml NGF and in the presence of FK506  
20 at 0.1 ng/ml NGF. At maximal concentrations of NGF (10-100 ng/ml), FK506 fails to produce additional neurite outgrowth.

FK506 is extremely potent in its neurotrophic effects. In the presence of a submaximal  
25 concentration of NGF (1 ng/ml) FK506 at 1 nM elicits the same maximal outgrowth observed with 50 ng/ml NGF (Fig. 11). Half maximal effects of FK506 occur at approximately 100 pM. In the absence of NGF,

FK506 fails to elicit neurite outgrowth (Fig. 10).

Rapamycin is a potent immunosuppressant which is not thought to act through calcineurin but which may influence other phosphorylation cascades.

5 Rapamycin potently blocks actions of FK506 that occur through FKBP and calcineurin presumably by acting as an FK506 antagonist at FKBP. Rapamycin (1  $\mu$ M) fails to block the neurotrophic actions of FK506. Instead, rapamycin is itself neurotrophic  
10 providing major neurite outgrowth at 1 nM. Rapamycin and FK506 seem to be acting via different mechanisms. Thus, rapamycin augments the number of processes as well as their length, while FK506 primarily increases neurite length. Moreover,  
15 effects of FK506 and rapamycin appear to be additive.

#### Dorsal Root Ganglia

This example demonstrates that FK506 is neurotrophic for sensory ganglia. We examined the  
20 action of FK506 on primary cultures of dorsal root ganglia from rats at embryonic day 16.

Stage E16 embryos were removed from pregnant Sprague-Dawley rats and the dorsal root ganglia dissected. Whole ganglia explants were cultured in  
25 collagen-coated 35 mm dishes (Falcon) using N2 medium (Dulbecco's Modified Eagle medium and Ham's F12 medium mixed 1:1 and supplemented with progesterone, selenium, insulin, putrescine,

glucose, and penicillin-streptomycin) at 37° C in a 15% CO<sub>2</sub> environment. Sensory ganglia were treated with various concentrations of NGF and/or FK506 or rapamycin or anti-NGF antibody. Ganglia were  
5 observed every 2-3 days under phase-contrasting using an Olympus IMT-2 inverted microscope, and measurements of axon length were made. The axonal field of each ganglion was divided into four quadrants, and the length of the longest axons in  
10 each quadrant was measured in microns using an eye-piece micrometer. The average of these measurements was taken as the axon length for the ganglion.

For [<sup>3</sup>H]FK506 autoradiography, dorsal root ganglia cultures were grown on chamber slides coated  
15 with collagen, 5 µg/cm<sup>2</sup>. Cultures were fixed on the slide with ice cold 4.0% freshly depolymerized paraformaldehyde in 0.1 M sodium phosphate buffer, pH 7.4, for 1 hr, then washed two times with phosphate buffered saline. Fixed cultures were  
20 labeled with [<sup>3</sup>H]FK506 by pre-incubating the slides in a buffer consisting of 50 mM Hepes, 2 mg/ml bovine serum albumin, 0.02% Tween-20 pH 7.4. This was followed by incubation in the same assay buffer containing 1 nM [<sup>3</sup>H]FK506. Non-specific binding was  
25 determined by adding 1 µM unlabeled FK506. The slides were then rinsed 4 x 5 min prior to drying, and juxtaposed to tritium-sensitive film for 10 days.

Autoradiography of [<sup>3</sup>H]FK506 binding sites reveals substantial levels of FKBP associated silver grains in these ganglia (Fig. 12). At 1  $\mu$ M unlabeled FK506, autoradiographic grains are

5 abolished indicating the specificity of binding. As reported previously, NGF (100 ng/ml) markedly increases the number and length of ganglia processes (Fig. 13). FK506 (1  $\mu$ M) alone produces a similar neurotrophic effect, while as little as 1 nM FK506

10 produces a noticeable increase in growth. Rapamycin (1  $\mu$ M) which acts as an FK506 antagonist, completely blocks the effects of FK506 (1  $\mu$ M), thus the action of FK506 displays a drug specificity characteristic of FKBP.

15       Whereas FK506 fails to stimulate neurite outgrowth in PC-12 cells in the absence of added NGF, in sensory ganglia FK506 alone is neurotrophic. Schwann cells in the ganglia can fabricate NGF, and the production of NGF by Schwann cells is regulated

20 by a protein phosphorylation event. To ascertain whether the actions of FK506 alone involve potentiation of endogenous NGF, we examined the influence of antibodies to NGF (Fig. 13). Anti-NGF markedly reduces the neurotrophic effects of FK506

25 (1  $\mu$ M). The anti-NGF is not acting in a toxic fashion as we observe no morphologic evidence of toxicity in the cells exposed to anti-NGF in the presence or absence of added NGF.

FK506 is extremely potent in stimulating neurite outgrowth. As little as 1 pM FK506 produces detectable augmentation. Progressively greater outgrowth occurs at 0.1 and 10 nM FK506 (data not shown), while maximal outgrowth requires 1  $\mu$ M FK506.

The time course of neurite outgrowth is similar at all concentrations of NGF and FK506. Some outgrowth is evidence by 1 day, while growth begins to plateau at about 5-6 days.

FK506 neurotrophic effects involve FKBP (FK506 binding protein) in sensory ganglia since the effects of FK506 are reversed by low concentrations of rapamycin, a known antagonist of FK506 at FKBP. The failure of rapamycin to block FK506 effects in PC-12 cells probably reflects the separate stimulatory effects of rapamycin. Mechanisms for rapamycin stimulation of neurite outgrowth in PC-12 cells are not immediately evident. Its immunosuppressant actions are thought to involve different mechanisms than those of FK506. Rapamycin can inhibit S6 kinase which phosphorylates the S6 ribosomal subunit. Rapamycin also inhibits phosphatidylinositol-3-kinase.

Protein kinase C (PKC) -mediated phosphorylation has been implicated in process outgrowth during neuronal regeneration. Other evidence suggests inhibitory effects of PKC in neuronal process extension.

GAP43 is a prominent calcineurin substrate highly concentrated in neurites and its phosphorylation is regulated by FKBP. GAP43 may not be directly involved in neurite extension, as PC-12 cell lines with low levels of GAP43 display normal neurite outgrowth. However, GAP43 and its phosphorylation may be involved in targeting neurites, as levels of phosphorylated GAP43 are increased when neurites approach their targets.

Phosphorylation of GAP-43 may also influence mobilization of  $\text{Ca}^{2+}$  that regulates neurite extension. Phosphorylated GAP-43 inhibits phosphatidyl inositol bis-phosphate formation, which should diminish level of inositol 1,4,5-triphosphate and associated  $\text{Ca}^{2+}$  release. In addition, phosphorylation of GAP-43 decreases its affinity for calmodulin with the resultant free calmodulin available to bind  $\text{Ca}^{2+}$ .

Immunophilins may act at sites besides calcineurin which affect  $\text{Ca}^{2+}$  that regulates neurite outgrowth. FKBP binds to the ryanodine receptor, which is a  $\text{Ca}^{2+}$  release channel. In skeletal muscle sarcoplasmic reticulum FK506 dissociates FKBP from the ryanodine receptor to facilitate the  $\text{Ca}^{2+}$  induced  $\text{Ca}^{2+}$  release mechanism. In addition, FK506 acts at other sites including FKBP25 steroid receptors and other unidentified targets such as those related to FKBP13. Thus other potential mechanisms may play some role in neurite extension.

Non-Immunosuppressive and Immunosuppressive Ligands  
of Immunophilins Stimulate Neurite Outgrowth in PC-  
12 Cells

In the present study we have examined in detail  
5 influences of a number of ligands of the  
immunophilins upon neurite extension in PC-12 cells  
and in intact chick sensory ganglia. We report that  
non-immunosuppressive as well as immunosuppressive  
ligands are extremely potent in augmenting neurite  
10 outgrowth in both PC-12 cells and sensory ganglia.

In our earlier study we found that  
immunosuppressants stimulate neurite outgrowth in  
PC-12 cells by increasing the potency of nerve  
growth factor (NGF) about 10-fold (Lyons et. al.,  
15 1994). In the absence of added NGF no neurotrophic  
effects are observed. In the present study we  
evaluated effects of the immunosuppressant drugs  
FK506 and rapamycin on PC-12 neurite outgrowth in  
the presence of 0.1 - 100 ng/ml of NGF.

20 In the absence of added NGF, none of the drugs  
stimulate neurite outgrowth. At 0.1 ng/ml NGF alone  
produces a small increase in neurite extension only  
about 15% of maximal effects that are evident at  
50 ng/ml (Figure 14). Rapamycin stimulates neurite  
25 outgrowth to a greater extent than the other drugs,  
with a 3-4 fold stimulation at 0.1 - 0.5 ng/ml NGF.  
The extent of augmentation elicited by rapamycin  
decreases with higher concentration of NGF and is

not statistically significant at 5-50 ng/ml NGF.  
FK506 also is neurotrophic with effects most  
apparent at lower NGF concentrations and a maximal  
2.5-fold enhancement of neurite outgrowth evident at  
5 0.5 ng/ml NGF.

There are three principal structural classes of  
immunosuppressant drugs related in structure to  
cyclosporin A, FK506, and rapamycin. Though FK506  
and cyclosporin A bind to distinct immunophilin  
10 proteins, they both act as immunosuppressants by  
inhibiting calcineurin. Rapamycin bind with very  
high affinity to FKBP-12, but the drug-immunophilin  
complex does not in turn bind to calcineurin.  
Instead, immunosuppressant actions result from the  
15 rapamycin-FKBP-12 complex binding to a recently  
identified and cloned protein designated RAFT-1  
(rapamycin and FK506 target) and also designated  
FRAP (Sabatini and Snyder, 1994; Brown et. al.,  
1994; Chen et. al., 1994). Because rapamycin binds  
20 potently to FKBP-12 but does not inhibit  
calcineurin, it can serve as an antagonist to FK506.  
There exist non-immunosuppressive derivatives of  
rapamycin. One of these, WAY-124,466, a triene  
derivative of rapamycin, binds with high affinity to  
25 FKBP-12 and inhibits rotamase activity, but is  
devoid of immunosuppressant actions. Cyclosporin A  
is a large cyclic undecapeptide. The mere addition  
of a methyl group to an alanine at the 6 position



results in an agent that does not inhibit calcineurin and lacks immunosuppressive effects, though it inhibits the rotamase activity of cyclophilin to a similar extent as cyclosporin A (Me  
5 CsA ref).

To ascertain whether immunosuppressant activity is required for neurotrophic actions, we compared the neurotrophic effects of FK506, rapamycin and cyclosporin A with non-immunosuppressant  
10 WAY-124,466, evaluating a wide range of concentrations on PC-12 cells (Figs. 15, 16). All studies were done in the presence of 0.5 ng/ml NGF. As observed previously, FK506 very potently stimulates neurite extension with half-maximal  
15 stimulation at 0.5 nM and maximal effects at 5-100 nM.

Rapamycin is the most potent agent examined and produces the greatest maximal level of neurite extension. In repeated experiments 50% of maximal  
20 extension is evident at about 0.2-0.4 nM while maximal effects are evident at about 10-100 nM. Maximal neurite extension with rapamycin is comparable to maximal effects of 50 ng/ml NGF. WAY-124466 also is neurotrophic, but is less potent and  
25 produces a lesser maximal effect than rapamycin. Half-maximal stimulation with WAY-124,466 occurs at about 10 nM and maximal effects occur at 100-1,000 nM. Thus, rapamycin is about 100-fold more potent

than WAY-124,466, resembling its 40-fold greater potency in binding to FKBP-12 (Table II).

Cyclosporin A is substantially less potent than FK506 or rapamycin in stimulating neurite outgrowth corresponding to its substantially lesser potency inhibiting rotamase activity. Fifty percent maximal stimulation of neurite outgrowth with cyclosporin is evident at 50 nM with maximal effects at 100 nM and a decrease in neurite outgrowth at higher concentrations of cyclosporin A. Maximal stimulation with cyclosporin A is about 60% of effects of 50 ng/ml NGF.

The general pattern of process extension is similar with the various immunophilin ligands and with NGF. At concentrations that elicit 50% of maximal effects, NGF (1-5 ng/ml) 40-50% of cells extend processes at least as long as the cell body while 15% extend longer processes, up to 3-5 times the length of the cell body. The pattern is fairly similar with the various drugs examined. Rapamycin and WAY-124,466 tend to result in a greater number of processes per cell than FK506. Cyclosporin A tends to be intermediate in terms of numbers of processes.

**Nerve Extension Elicited in Chick Dorsal Root Ganglia by Non-Immunosuppressive and Immunosuppressive Ligands of Immunophilins**

In our previous study, we observed neurite extension

effects of immunosuppressant drugs in explants of rat dorsal root ganglia with significant augmentation in nerve outgrowth observed with FK506 concentrations as low as 1 picomolar (Lyons et. al., 1994). In the rat ganglia neurotrophic effects were observed with FK506 even in the absence of NGF. In the present study we have employed explants of chick dorsal root ganglia, which are easier to employ in studies of nerve outgrowth. In the absence of added NGF, we observe minimal effects of immunophilin ligand drugs. The chick cells are more sensitive to NGF than PC-12 cells so that we employ 0.1 ng/ml NGF to produce minimal neurite outgrowth and to demonstrate neurotrophic actions of immunophilin ligands (Fig. 17,18).

Dorsal root ganglion were dissected from chick embryos of ten day gestation. Whole ganaglion explants were cultured on thin layer Matrigel-coated 12 well plates with Liebovitz L15 plus high glucose media supplemented with 2mM glutamine and 10% fetal calf serum, and also containing 10  $\mu$ M cytosine  $\beta$ -D arabinofuranoside (Ara C) at 37°C in an environment containing 5% CO<sub>2</sub>. Twenty-four hours later, the DRG's were treated with various conentrations of nerve growth factor, immunophilin ligands or combinations of NGF plus drugs. Forty-eight hours after drug treatment, the ganglia were visualized under phase contrast or Hoffman Modulation contrast

with a Zeiss Axiovert inverted microscope.

Photomicrographs of the explants were made, and neurite outgrowth was quantitated. Neurites longer than the DRG diameter were counted as

5 positive, with total number of neurites quantitated per each experimental condition. Three to four DRGs are cultured per well, and each treatment was performed in duplicate.

The relative potencies of the various  
10 immunophilin ligands in stimulating nerve outgrowth in the ganglia are similar to their relative potencies in PC-12 cells. Thus, rapamycin is the most potent agent with an  $EC_{50}$  of 1nM, 10-fold more potent than WAY-124,466, while FK506 displays  
15 an  $EC_{50}$  of 1-2nM.

The maximal increase in the number of processes, their length and branching is quite similar at maximally effective concentrations of the immunophilin ligands and of NGF (100 ng/ml). With  
20 progressively increasing concentrations of the various drugs, one observes a larger number of processes, more extensive branching and a greater length of individual processes.

We evaluated the potencies of drugs in binding  
25 to FKBP-12 by examining inhibition of  $^3H$ -FK506 binding to recombinant FKBP-12. There is a striking parallel between affinities of drugs for FKBP-12 and their potencies in stimulating neurite outgrowth and

inhibiting rotamase activity. Clearly, stimulation of nerve outgrowth is unrelated to calcineurin inhibition. Calcineurin inhibition fits well with immunosuppressant actions, WAY-124,466 is not immunosuppressive and fails to inhibit calcineurin. Rapamycin is a potent immunosuppressant, but the rapamycin-FKBP-12 complex binds to RAFT-1 to initiate immunosuppressive processes (Sabatini and Snyder, 1994; Snyder and Sabatini, 1995). The results are set forth in Table 2.

TABLE 2  
IMMUNOPHILIN LIGAND NEUROTROPHISM PARALLELS INHIBITION OF ROTAMASE, NOT CALCINEURIN

Drug	[ <sup>3</sup> H]-FK506-FKBP12 (IC <sub>50</sub> )	CALCINEURIN ROTAMASE INHIBITION	NEURITE OUTGROWTH (K <sub>T</sub> )	(ED <sub>50</sub> )
FK506	0.6 nM	YES 0.4 nM		0.5 nM
Rapamycin	0.5 nM	NO	0.2 nM	0.5 nM
WAY-124466	10.0 nM	NO	12.0 nM	10 nM
Cyclosporin A (CsA)	None	YES 20 nM		50 nM

We compared the ability of non-immunosuppressive immunophilin ligands to promote neurite outgrowth in chick dorsal root ganglion explant cultures (Table 3). Each of these compounds is incapable of inhibiting calcineurin, but they interact with the immunophilin FKBP-12 to inhibit its rotamase activity with the various inhibitory constants listed in Table 3. The ability of these compounds to promote neurite outgrowth in the DRG's correlates well with their ability to inhibit the rotamase activity of FKBP-12.

Table 3 IMMUNOPHILIN LIGAND NEUROTROPHISM PARALLELS INHIBITION  
OF ROTAMASE, NOT CALCINEURIN

<u>Drug</u>	<u>[<sup>3</sup>H]-FK506- FKBP12 (IC<sub>50</sub>)</u>	<u>Calcineurin Inhibition</u>	<u>Rotamase (K<sub>i</sub>)</u>	<u>Neurite Outgrowth (ED<sub>50</sub>)</u>
5 Example 12	8 $\mu$ M	NO	250 nM	300 nM
Example 13	4 $\mu$ M	NO	25 nM	80 nM

The very close correlation between the potencies of drugs in binding to immunophilins, inhibiting their rotamase activity and stimulating neurite outgrowth implies that inhibition of rotamase activity is responsible for neurotrophic effects of the drugs. The extraordinarily high potency of the drugs in stimulating neurite outgrowth and in binding to immunophilins makes it most unlikely that any other target could account for the neurotrophic effects. It is conceivable that a biological activity of immunophilins other than rotamase could be influenced by the drugs to mediate neurotrophic actions. However, no such activity has yet been reported.

Because of the extraordinary potency of the drugs and the close correlation between rotamase inhibition and neurotrophic actions, we conclude that rotamase inhibition is likely involved in neurotrophic effects. A number of proteins have been reported as substrates for the rotamase activity of immunophilins including collagen (Steinmann et. al., 1991) and transferrin (Lodish and King, 1991). Recently highly purified preparations of ryanodine receptor and the IP-3 receptor, prominent intracellular calcium channels have been reported to

exist in a complex with FKBP-12. Dissociation of FKBP-12 from these complexes causes the calcium channel to become "leaky" (Cameron et. al., 1995) Calcium fluxes are involved in neurite extension so that the IP-3 receptor and the 5 ryanodine receptor might be involved in the neurotrophic effects of drugs. Since the drugs bind to the same site on FKBP-12 as the IP-3 receptor or the ryanodine receptor, one would have to postulate that the drugs displace the channels from FKBP-12. No interaction between these calcium channels 10 in cyclophilin has been reported so that this model would not explain the neurotrophic actions of cyclosporin A.


The neurotrophic actions of the drugs studied here are exerted at extremely low concentrations indicating potencies comparable to those of neurotrophic proteins such as brain 15 derived growth factor, neurotrophin-3 and neurotrophic growth factor.

The following examples are illustrative of preferred embodiments of the invention and are not to be construed as limiting the invention thereto. All polymer molecular 20 weights are mean average molecular weights. All percentages are based on the percent by weight of the final delivery system or formulation prepared unless otherwise indicated and all totals equal 100% by weight.

Illustrative pipecolic acid derivative compounds which 25 can be used for the purposes of this invention include:

46


## EXAMPLE 1



Way-124,466

This exemplary pipecolic acid derivative compound is disclosed by Ocain et al., *Biochemical and Biophysical Research Communications*, Vol. 192, No. 3, 1993. The compound was synthesized at Wyeth-Ayerst by Dr. Phil Hughes by reaction of 4-phenyl-1,2,4-triazoline-3,5-dione with rapamycin.

## EXAMPLE 2




RAP-Pa



47

This pipecolic acid derivative compound is disclosed by Chakraborty et al., *Chemistry and Biology*, March 1995, 2:157-161.

## EXAMPLES 3-5




5 Exemplary pipecolic acid derivative compounds are disclosed by Ikeda et al., *J. Am. Chem. Soc.* 1994, 116, 4143-4144, and are incorporated herein by reference.

## EXAMPLES 6-9

10 Exemplary pipecolic acid derivative compounds are disclosed by Wang et al., *Bioorganic and Medicinal Chemistry Letters*, Vol. 4, No. 9, pp. 1161-1166, 1994, particularly compounds 2a-2d and are incorporated herein by reference.

## EXAMPLE 10




This exemplary pipecolic acid derivative, compound 10, is disclosed by Birkenshaw et al., *Bioorganic & Medicinal Chemistry Letters*, Vol. 4, No. 21, pp. 2501-2506, 1994, and is incorporated herein by reference.

## EXAMPLES 11-21

Exemplary pipecolic acid derivative compounds are disclosed by Holt et al., *J. Am. Chem. Soc.*, 1993, 115, 9925-9938, particularly compounds 4-14, and are incorporated herein by reference.

## EXAMPLES 22-30

Exemplary pipecolic acid derivative compounds are disclosed by Caffery et al., *Bioorganic & Medicinal Chemistry Letters*, Vol. 4, No. 21, pp. 2507-2510, 1994, and are incorporated herein by reference.

EXAMPLE 31

This exemplary pipecolic acid derivative, compound 31, is disclosed by Teague et al., *Bioorganic & Medicinal Chemistry Letters*, Vol. 3, No. 10, pp. 1947-1950, 1993 and is incorporated herein by reference.

EXAMPLES 32-34

Exemplary pipecolic acid derivative compounds are disclosed by Yamashita et al., *Bioorganic & Medicinal Chemistry Letters*, Vol. 4., No. 2, pp. 325-328, 1994, particularly, compounds 11, 12, and 19, and are incorporated herein by reference.

EXAMPLE 35-55

Exemplary pipecolic acid derivatives are disclosed by Holt et al., *Bioorganic & Medicinal Chemistry Letters*, Vol. 4, No. 2, pp. 315-320, 1994, particularly, compounds 3-21, and 23-24, and are incorporated herein by reference.


## EXAMPLES 56-68

Exemplary pipecolic acid derivative compounds are disclosed by Holt et al., *Bioorganic & Medicinal Chemistry Letters*, Vol. 3, No. 10, pp. 1977-1980, 1993, particularly compounds 3-15 and are incorporated by reference herein.

## EXAMPLES 69-83

Exemplary compounds of the present invention are disclosed by Hauske et al., *J. Med. Chem.* 1992, 35, 4284-4296, particularly compounds 6, 9-10, 21-24, 26, 28, 31-32, and 52-55, and are incorporated herein by reference.

## EXAMPLE 84



## SLB506

This exemplary pipecolic acid derivative is disclosed by Teague et al., *Bioorganic & Med. Chem. Letters*, Vol. 4, No. 13, pp. 1581-1584, 1994, and is incorporated herein by reference.


51

**EXAMPLES 85-88**

Exemplary pipecolic acid derivative compounds are disclosed by Stocks et al., *Bioorganic & Med. Chem. Letters*, Vol. 4, No. 12, pp. 1457-1460, 1994, particularly compounds 2, 15-17 and are incorporated herein by reference.


**EXAMPLES 90-111**

Additional exemplary pipecolic acid derivatives are described in Scheme 10, Tables 1-5.

**SCHEME 01**

EXAMPLE/COMPOUND	STRUCTURE
6	X = H,
7	X = CH <sub>2</sub> ,
8	X = H, CH <sub>2</sub> ,
9	X = O

## SCHEME 2



## EXAMPLE/COMPOUNDS No.

R<sub>2</sub>


	11	
	12	
5	13	
	14	
	15	
	16	
	17	
10	18	

## SCHEME 3


EXAMPLE/COMPOUND No.

STRUCTURE

19




20




5

21



Scheme 4


Table 1



5

Example/Compound No.	Structure
24	y=1
23	y=2
24	y=3

Table 2




10

Example/Compound No.	Structure
25	n=1
26	n=2
27	n=3




Table 3.



Example/Compound No.	Structure
28	n=1
29	n=2
30	n=3

SCHEME 5



EXAMPLE/COMPOUND No.

STRUCTURE

32


R=phenyl

33

R=N(allyl)<sub>2</sub>

10


34



56

Scheme 6

Table 1




Example/Compound No.	Structure
R	
5	35
	36
	37
10	38
	39
	40
15	41
	42
	43
20	
	44
	45
20	46
	47
	48
20	
	49
20	50

Table 2


Example/Compound No.

Structure


51




52




53



54




55



53


Scheme 7

TABLE 1



5	56	x = OH
	57	x = OMe
	58	x = Oi Pr
	59	x = OBn
	60	x = OCH <sub>2</sub> MePh
10	61	x = OCH <sub>2</sub> CHCHPh
	62	x = OCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> (3,4-OMe <sub>2</sub> )Ph
	63	x = NHBn
	64	x = NHCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> Ph

Table 2




65	R = Me
66	R = Bn


55

Table 3

57




68



## Scheme 8


Table 1



Example/Compound No.

Structure

69

 $n = 2, R_1 =$  $R_2 = \text{Phe-o-tert-butyl}$ 

5

70



 $n = 2, R_1 =$  $R_2 = \text{Phe-o-tert-butyl}$ 

Table 2



10

71

 $R_1 = m\text{-OCH}_3\text{Ph}; R_2 = \text{Val-o-tert-butyl}$ 

72

 $R_1 = m\text{-OCH}_3\text{Ph}; R_2 = \text{Leu-o-tert-butyl}$ 

73

 $R_1 = m\text{-OCH}_3\text{Ph}; R_2 = \text{Ileu-o-tert-butyl}$ 

74

 $R_1 = m\text{-OCH}_3\text{Ph}; R_2 = \text{hexahydro-Phe-o-tert-butyl}$ 

75

 $R_1 = m\text{-OCH}_3\text{Ph}; R_2 = \text{allylalanine-o-tert-butyl}$ 


15

76

 $R_1 = 3\text{-naphthyl}; R_2 = \text{Val-o-tert-butyl}$

61

Table 3



Example/Compound No.

Structure

77

 $R_1 = \text{CH}_2(\text{CO})-m\text{-OCH}_3\text{Ph}$  $R_4 = \text{CH}_2\text{Ph}$  $R_5 = \text{OCH}_3$ 


5

78

 $R_1 = \text{CH}_2(\text{CO})-\beta\text{-naphthyl}$  $R_4 = \text{CH}_2\text{Ph}$  $R_5 = \text{OCH}_3$

62

Table 4



Example/Compound No.	Structure
5	79 $R_1 = m\text{-OCH}_3\text{Ph}$ $X = \text{trans-CH=CH}$ $R_{4,1} = \text{H}$ $Y = \text{OC(o)Ph}$
10	80 $R_1 = m\text{-OCH}_3\text{Ph}$ $X = \text{trans-CH=CH}$ $R_{4,1} = \text{H}$ $Y = \text{OC(o)CF}_3$
15	81 $R_1 = m\text{-OCH}_3\text{Ph}$ $X = \text{trans-CH=CHI}$ $R_{4,1} = -$ $Y = -$
15	82 $R_1 = m\text{-OCH}_3\text{Ph}$ $X = \text{trans-CH=CH}$ $R_{4,1} = \text{H}$ $Y = \text{OCH}_2\text{CH=CH}_2$
20	83 $R_1 = m\text{-OCH}_3\text{Ph}$ $X = \text{C=O}$ $R_{4,1} = \text{H}$ $Y = \text{Ph}$



Scheme 9  
Table 1  
85





Table 2




86  
87  
88

$R_1 = H, R_2 = OMe, R_3 = CH_2OMe$   
 $R_1 = H, R_2 = R_3 = H$   
 $R_1 = Me, R_2 = R_3 = H$


## Scheme 10

Table 1



	Example	R =
5	90	3,4-dichloro
	91	3,4,5-trimethoxy
	92	H
	93	3-(2,5-Dimethoxy) - phenylpropyl
10	94	3-(3,4-Methylene- dioxy) phenylpropyl


Table 2



	Example	R =
15	95	4-(p-Methoxy) -butyl
	96	3-Phenylpropyl
	97	3-(3-Pyridyl) - propyl


65

Table 3




	Example	R =
5	98	3-(3-Pyridyl)-propyl
	99	1,7-Diphenyl-4-heptyl
	100	4-(4-Methoxy)butyl
	101	1-Phenyl-6-(4-methoxy-phenyl)-4-hexyl
	102	3-(2,5-Dimethoxy)phenyl-propyl
10	103	3-(3,4-Methylenedioxy)-phenylpropyl
	104	1,5-Diphenylpentyl

Table 4



	Example	R =
15	105	4-(4-Methoxy)butyl
	106	3-Cyclohexylpropyl
	107	3-Phenylpropyl

Table 5



20

	Example	R =
25	108	3-Cyclohexylpropyl
	109	3-Phenylpropyl
	110	4-(4-Methoxy)butyl
	111	1,7-Diphenyl-4-
		heptyl

## NEUROTROPHIC EFFECTS OF ROTAMASE INHIBITORS

Table I lists a number of the claimed examples together with their potencies to induce trophic effects in cultured sensory neurons, as described above.

5 Figures 19 and 20 show photomicrographs of Example 111 and Example 17 promoting neurite outgrowth in the dorsal root ganglion cultures.

Table I  
In Vitro Potencies of Test Examples

10	Example	Rotamase Inhibition K <sub>i</sub> , nM	Neutrophic ED <sub>50</sub> Chick DRGs, nM
	6	140	25
	9	13	0.030
	11	170	1
15	12	250	300
	13	25	80
	15	17	0.30
	19	12	0.017
	36	>10,000	>10,000
20	41	1300	5000
	50	>10,000	>10,000
	90	1800	2500
	91	28	200
	92	39	90
25	93	75	35
	94	70	8
	95	165	5-10
	96	740	10-20
	97	725	150
30	98	130	75
	99	30	5
	100	60	43
	101	15	0.17
	102	12	2.5
35	103	120	3
	104	20	.016
	105	103	6
	106	760	1
	107	210	0.82
40	108	32	0.29
	109	2	0.08
	110	24	0.002
	111	5	0.08

ACTIVITY OF EXAMPLE COMPOUNDS IN IN VIVO MODEL OF  
NERVE REGENERATION

Sciatic Nerve Axotomy

5       Six-week old male Sprague-Dawley rats were  
anesthetized, and the sciatic nerve exposed and crushed,  
at the level of the hip, by forceps. Test compounds or  
vehicle were administered subcutaneously just prior to  
the lesion and daily for the following 18 days. Sections  
of the sciatic nerve were stained with Holmes silver  
10       stain to quantify the number of axons, and Luxol fast  
blue to quantify the level of myelination. Eighteen days  
after lesion, there was a significant decrease in the  
number of axons (50% decrease as compared to non-lesioned  
control) and degree of myelination (90% decrease as  
15       compared to non-lesioned control) in animal treated with  
vehicle.

Administration of Example 12 (30 mg/kg s.c.), or  
Example 13 (mg/kg s.c.) just prior to the lesion and  
daily for 18 days following the lesion, resulted in  
20       significant regeneration of both axon number (25% and 5%  
decrease, respectively, as compared to non-lesioned  
control) and the degree of myelination (65 % and 50%  
decrease, respectively, as compared to control) as  
compared to vehicle treated animals. The significant  
25       efficacy of Examples 12 and 13 are consistent with their  
potent activity in inhibiting rotamase activity and  
stimulating neurite outgrowth in chick DRGs, and their  
relative potencies in vivo parallel their in vitro  
potencies (Table I). These results are shown in Figure  
30       21. "Sham" denotes control animals that received vehicle  
but were not lesioned; "Vehicle" denotes animals that  
were lesioned and received only vehicle (i.e., no drug).  
Example 12 and Example 13 showed a striking similarity to  
the sham treated animals, demonstrating the powerful  
35       neuroregenerative effects of these compounds in vivo.  
These data are quantitated in Table II.

68

Table II

Treatment	Axon Number (% Control)	Myelin Level
Sham	100	100
5 Lesion:		
+ Vehicle (s.c.)	50	10
+ Example 12 (30 mg/kg s.c.)	75	35
10 + Example 13 (30 mg/kg s.c.)	100	50

The invention being thus described, it will be obvious that the same may be varied in many ways. Such variations are not to be regarded as a departure from the spirit and scope of the invention and all such modifications are intended to be included within the scope of the following claims.

What is claimed is:

1. A method of treating a neurological disorder in an animal, comprising:  
administering to an animal an effective amount of a  
5 pipecolic acid derivative having an affinity for FKBP-type immunophilins to stimulate growth of damaged peripheral nerves or to promote neuronal regeneration, wherein the FKBP-type immunophilin exhibits rotamase activity and the pipecolic acid derivative inhibits said rotamase activity of the  
10 immunophilin.
2. The method of claim 1, wherein the neurological disorder is selected from the group consisting of peripheral neuropathies cause by physical injury or disease state, physical damage to the brain, physical damage to the spinal  
15 cord, stroke associated with brain damage, and neurological disorders relating to neurodegeneration.
3. The method of claim 2, wherein the neurological disorder is selected from the group consisting of Alzheimer's Disease, Parkinson's Disease, and amyotrophic  
20 lateral sclerosis.
4. The method of claim 1, wherein the pipecolic acid derivative compound is immunosuppressive or non-immunosuppressive.
5. The method of claim 1, wherein the pipecolic acid  
25 derivative is Way-124,666.
6. The method of claim 1, wherein the pipecolic acid derivative is rapamycin.
7. The method of claim 1, wherein the pipecolic acid derivative is FK506.
- 30 8. The method of claim 1, wherein the pipecolic acid derivative is Rap-Pa.

9. The method of claim 1, wherein the pipecolic acid derivative is SLB-506.
10. The method of claim 1, wherein the pipecolic acid derivative is selected from the group consisting of compounds  
5 3-84, and 86-88.
11. A method of treating a neurological disorder in an animal, comprising:  
administering to an animal an effective amount of a pipecolic acid derivative having an affinity for FKBP-type  
10 immunophilins in combination with an effective amount of a neurotrophic factor selected from the group consisting of neurotrophic growth factor, brain derived growth factor, glial derived growth factor, ciliary neurotrophic factor, and neurotrophin-3, to stimulate growth of damaged peripheral  
15 nerves or to promote neuronal regeneration, wherein the FKBP-type immunophilin exhibits rotamase activity and the pipecolic acid derivative inhibits said rotamase activity of the immunophilin.
12. The method of claim 11, wherein the neurological  
20 disorder is selected from the group consisting of peripheral neuropathies caused by physical injury or disease state, physical damage to the brain, physical damage to the spinal cord, stroke associated with brain damage, and neurological disorders relating to neurodegeneration.
- 25 13. The method of claim 12, wherein the neurological disorder is selected from the group consisting of Alzheimer's Disease, Parkinson's Disease, and amyotrophic lateral sclerosis.
- 30 14. The method of claim 11, wherein the pipecolic acid derivative compound is immunosuppressive or non-immunosuppressive.



15. The method of claim 11, wherein the pipecolic acid derivative is Way-124,666.
16. The method of claim 11, wherein the pipecolic acid derivative is rapamycin.
- 5 17. The method of claim 11, wherein the pipecolic acid derivative is FK506.
18. The method of claim 11, wherein the pipecolic acid derivative is Rap-Pa.
19. The method of claim 11, wherein the pipecolic acid  
10 derivative is SLB-506.
20. The method of claim 11, wherein the pipecolic acid derivative is selected from the group consisting of compounds 3-84, and 86-88.
21. A method of stimulating growth of damaged peripheral  
15 nerves, comprising;  
administering to damaged peripheral nerves an effective amount of a pipecolic acid derivative compound having an affinity for FKBP-type immunophilins to stimulate or promote growth of the damaged peripheral nerves, wherein the FKBP-  
20 type immunophilins exhibit rotamase activity and the pipecolic acid derivative inhibits said rotamase activity of the immunophilin.
22. The method of claim 21, further comprising  
administering a neurotrophic factor to stimulate or promote  
25 growth of the damaged peripheral nerves selected from the group consisting of neurotrophic growth factor, brain derived growth factor, glial derived growth factor, ciliary neurotrophic factor, and neurotrophin-3.
23. The method of claim 21, wherein the pipecolic acid  
30 derivative is immunosuppressive or non-immunosuppressive.

24. The method of claim 21, wherein the pipecolic acid derivative is Way-124,666

25. The method of claim 21, wherein the pipecolic acid derivative is rapamycin.

26. The method of claim 21, wherein the pipecolic acid derivative is FK506.

27. The method of claim 21, wherein the pipecolic acid derivative is Rap-Pa.

28. The method of claim 21, wherein the pipecolic acid derivative is SLB-506.

29. The method of claim 21, wherein the pipecolic acid derivative is selected from the group consisting of compounds 3-84, and 86-88.

30. A method of stimulating growth of damaged peripheral nerves, comprising:

administering to damaged peripheral nerves an effective amount of a pipecolic acid derivative compound having an affinity for FKBP-type immunophilins to stimulate growth of damaged peripheral nerves, wherein the FKBP-type immunophilin exhibit rotamase activity and the pipecolic acid derivative inhibits said rotamase activity of the immunophilin.

31. The method of claim 30, further comprising administering an effective amount of a neurotrophic factor selected from the group consisting of neurotrophic growth factor, brain derived growth factor, glial derived growth factor, ciliary neurotrophic factor, and neurotrophin-3 to stimulate the growth of damaged peripheral nerves.

32. The method of claim 30, wherein the pipecolic acid derivative compound is immunosuppressive or non-immunosuppressive.

33. The method of claim 30, wherein the pipecolic acid derivative is Way-124,666
34. The method of claim 30, wherein the pipecolic acid derivative is rapamycin.
35. The method of claim 30, wherein the pipecolic acid derivative is FK506.
36. The method of claim 30, wherein the pipecolic acid derivative is Rap-Pa.
37. The method of claim 30, wherein the pipecolic acid derivative is SLB-506.
38. The method of claim 30, wherein the pipecolic acid derivative is selected from the group consisting of compounds 3-84, and 86-88.
39. A method for promoting neuronal regeneration and growth in animals, comprising:
  - administering to an animal an effective amount of a pipecolic acid derivative compound having an affinity for FKBP-type immunophilins to promote neuronal regeneration, wherein the FKBP-type immunophilins exhibit rotamase activity and the pipecolic acid derivative inhibits said rotamase activity of the immunophilin.
40. The method of claim 39, further comprising administering an effective amount of a neurotrophic factor to promote neuronal regeneration selected from the group consisting of neurotrophic growth factor, brain derived growth factor, glial derived growth factor, and neurotrophin-3.
41. The method of claim 39, wherein the pipecolic acid derivative compound is immunosuppressive or non-immunosuppressive.

42. The method of claim 39, wherein the pipecolic acid derivative is Way-124,666.

43. The method of claim 39, wherein the pipecolic acid derivative is rapamycin.

5 44. The method of claim 39, wherein the pipecolic acid derivative is FK506.

45. The method of claim 39, wherein the pipecolic acid derivative is Rap-Pa.

10 46. The method of claim 39, wherein the pipecolic acid derivative is SLB-506.

47. The method of claim 39, wherein the pipecolic acid derivative is selected from the group consisting of compounds 3-84, and 86-88.

15 48. A method for preventing neurodegeneration in an animal, comprising:

administering to an animal an effective amount of a pipecolic acid derivative having an affinity for FKBP-type immunophilins to prevent neurodegeneration, wherein the FKBP-type immunophilin exhibits rotamase activity and the pipecolic acid derivative inhibits said rotamas activity of the immunophilin.

20 49. The method of claim 48, further comprising administering an effective amount of a neurotrophic factor to prevent neurodegeneration selected from the group consisting of neurotrophic growth factor, brain derived growth factor, glial derived growth factor, ciliary neurotrophic factor, and neurotrophin-3.

25 50. The method of claim 48, wherein the pipecolic acid derivative compound is immunosuppressive or non-immunosuppressive.

51. The method of claim 48, wherein the pipecolic acid derivative is Way-124,666.
52. The method of claim 48, wherein the pipecolic acid derivative is rapamycin.
- 5 53. The method of claim 48, wherein the pipecolic acid derivative is FK506.
54. The method of claim 48, wherein the pipecolic acid derivative is Rap-Pa.
- 10 55. The method of claim 48, wherein the pipecolic acid derivative is SLB-506.
56. The method of claim 48, wherein the pipecolic acid derivative is selected from the group consisting of compounds 3-84, and 86-88.

## AMENDED CLAIMS

[received by the International Bureau on 28 November 1996 (28.11.96); original claims 4,6,7,14,16, 17,23,25,26,32,34,35,41,43,44,50,52 and 53 cancelled; original claims 1,5, 8-11,15,18-21,24,27-30,33,36-39,42,45-48, 51 and 54-56 amended; remaining claims unchanged (12 pages)]

1. A method of treating a neurological disorder in an animal, comprising:

administering to an animal an effective amount of a non-immunosuppressive pipecolic acid derivative having an affinity for FKBP-type immunophilins to stimulate growth of damaged peripheral nerves or to promote neuronal regeneration, wherein the FKBP-type immunophilin exhibits rotamase activity and the pipecolic acid derivative inhibits said rotamase activity of the immunophilin.

2. The method of claim 1, wherein the neurological disorder is selected from the group consisting of peripheral neuropathies cause by physical injury or disease state, physical damage to the brain, physical damage to the spinal cord, stroke associated with brain damage, and neurological disorders relating to neurodegeneration.

3. The method of claim 2, wherein the neurological disorder is selected from the group consisting of Alzheimer's Disease, Parkinson's Disease, and amyotrophic lateral sclerosis.

5. The method of claim 1, wherein the non-immunosuppressive pipecolic acid derivative is Way-124,666.

8. The method of claim 1, wherein the non-immunosuppressive pipecolic acid derivative is Rap-Pa.
9. The method of claim 1, wherein the non-immunosuppressive pipecolic acid derivative is SLB-506.
10. The method of claim 1, wherein the non-immunosuppressive pipecolic acid derivative is selected from the group consisting of compounds 3-84, 86-88 and 90-111.
11. The method of claim 1, wherein the method provides at least 75% regeneration of axon number and at least 35% recovery of myelination.
12. The method of claim 1, wherein the method provides at least 75% regeneration of axon number and at least 50% recovery of myelination.
13. The method of claim 1, wherein rotamase inhibition is achieved using no greater than a 1800 nM concentration of the compound.
14. The method of claim 1, wherein rotamase inhibition is achieved using no greater than a 210 nM concentration of the compound.
15. The method of claim 1, wherein rotamase inhibition is

achieved using no greater than a 103 nM concentration of the compound.

16. The method of claim 1, wherein neurotrophic  $ED_{50}$  is achieved using no greater than a 200 nM concentration of the compound.

17. The method of claim 1, wherein neurotrophic  $ED_{50}$  is achieved using no greater than a 35 nM concentration of the compound.

18. The method of claim 1, wherein neurotrophic  $ED_{50}$  is achieved using no greater than a 1 nM concentration of the compound.

19. A method of treating a neurological disorder in an animal, comprising:

administering to an animal an effective amount of a non-immunosuppressive pipecolic acid derivative having an affinity for FKBP-type immunophilins in combination with an effective amount of a neurotrophic factor selected from the group consisting of neurotrophic growth factor, brain derived growth factor, glial derived growth factor, ciliary neurotrophic factor, and neurotrophin-3, to stimulate growth of damaged peripheral nerves or to promote neuronal regeneration, wherein the FKBP-type immunophilin exhibits rotamase activity and the non-immunosuppressive pipecolic acid



derivative inhibits said rotamase activity of the immunophilin.

20. The method of claim 19, wherein the neurological disorder is selected from the group consisting of peripheral neuropathies cause by physical injury or disease state, physical damage to the brain, physical damage to the spinal cord, stroke associated with brain damage, and neurological disorders relating to neurodegeneration.

21. The method of claim 19, wherein the neurological disorder is selected from the group consisting of Alzheimer's Disease, Parkinson's Disease, and amyotrophic lateral sclerosis.

22. The method of claim 19, wherein the non-immunosuppressive pipecolic acid derivative is Way-124,666.

23. The method of claim 19, wherein the non-immunosuppressive pipecolic acid derivative is Rap-Pa.

24. The method of claim 19, wherein the non-immunosuppressive pipecolic acid derivative is SLB-506.

25. The method of claim 19, wherein the non-immunosuppressive pipecolic acid derivative is selected from the group consisting of compounds 3-84, 86-88 and 90-111.

26. A method of stimulating growth of damaged peripheral nerves, comprising;

administering to damaged peripheral nerves an effective amount of a non-immunosuppressive pipecolic acid derivative compound having an affinity for FKBP-type immunophilins to stimulate or promote growth of the damaged peripheral nerves, wherein the FKBP-type immunophilins exhibit rotamase activity and the non-immunosuppressive pipecolic acid derivative inhibits said rotamase activity of the immunophilin.

27. The method of claim 26, further comprising administering a neurotrophic factor to stimulate or promote growth of the damaged peripheral nerves selected from the group consisting of neurotrophic growth factor, brain derived growth factor, glial derived growth factor, ciliary neurotrophic factor, and neurotrophin-3.

28. The method of claim 26, wherein the non-immunosuppressive pipecolic acid derivative is Way-124,666

29. The method of claim 26, wherein the non-immunosuppressive pipecolic acid derivative is Rap-Pa.

30. The method of claim 26, wherein the non-immunosuppressive pipecolic acid derivative is SLB-506.

31. The method of claim 26, wherein the non-

immunosuppressive pipecolic acid derivative is selected from the group consisting of compounds 3-84, 86-88 and 90-111.

32. The method of claim 26, wherein the method provides at least 75% regeneration of axon number and at least 35% recovery of myelination.

33. The method of claim 26, wherein the method provides at least 75% regeneration of axon number and at least 50% recovery of myelination.

34. The method of claim 26, wherein rotamase inhibition is achieved using no greater than a 1800 nM concentration of the compound.

35. The method of claim 26, wherein rotamase inhibition is achieved using no greater than a 210 nM concentration of the compound.

36. The method of claim 26, wherein rotamase inhibition is achieved using no greater than a 103 nM concentration of the compound.

37. The method of claim 26, wherein neurotrophic  $ED_{50}$  is achieved using no greater than a 200 nM concentration of the compound.

38. The method of claim 26, wherein neurotrophic  $ED_{50}$  is achieved using no greater than a 35 nM concentration of the compound.

39. The method of claim 26, wherein neurotrophic  $ED_{50}$  is achieved using no greater than a 1 nM concentration of the compound.

40. A method of stimulating growth of damaged peripheral nerves, comprising:

administering to damaged peripheral nerves an effective amount of a non-immunosuppressive pipecolic acid derivative compound having an affinity for FKBP-type immunophilins to stimulate growth of damaged peripheral nerves, wherein the FKBP-type immunophilin exhibit rotamase activity and the non-immunosuppressive pipecolic acid derivative inhibits said rotamase activity of the immunophilin.

41. The method of claim 40, further comprising administering an effective amount of a neurotrophic factor selected from the group consisting of neurotrophic growth factor, brain derived growth factor, glial derived growth factor, ciliary neurotrophic factor, and neurotrophin-3 to stimulate the growth of damaged peripheral nerves.

42. The method of claim 40, wherein the non-immunosuppressive pipecolic acid derivative is Way-124,666.

43. The method of claim 40, wherein the non-immunosuppressive pipecolic acid derivative is Rap-Pa.

44. The method of claim 40, wherein the non-immunosuppressive pipecolic acid derivative is SLB-506.

45. The method of claim 40, wherein the non-immunosuppressive pipecolic acid derivative is selected from the group consisting of compounds 3-84, 86-88 and 90-111.

46. A method for promoting neuronal regeneration and growth in animals, comprising:

administering to an animal an effective amount of a non-immunosuppressive pipecolic acid derivative compound having an affinity for FKBP-type immunophilins to promote neuronal regeneration, wherein the FKBP-type immunophilins exhibit rotamase activity and the non-immunosuppressive pipecolic acid derivative inhibits said rotamase activity of the immunophilin.

47. The method of claim 46, further comprising administering an effective amount of a neurotrophic factor to promote neuronal regeneration selected from the group consisting of neurotrophic growth factor, brain derived growth factor, glial derived growth factor, and neurotrophin-3.

48. The method of claim 46, wherein the non-immunosuppressive pipecolic acid derivative is Way-124,666.
49. The method of claim 46, wherein the non-immunosuppressive pipecolic acid derivative is Rap-Pa.
50. The method of claim 46, wherein the non-immunosuppressive pipecolic acid derivative is SLB-506.
51. The method of claim 46, wherein the non-immunosuppressive pipecolic acid derivative is selected from the group consisting of compounds 3-84, 86-88 and 90-111.
52. The method of claim 46, wherein the method provides at least 75% regeneration of axon number and at least 35% recovery of myelination.
53. The method of claim 46, wherein the method provides at least 75% regeneration of axon number and at least 50% recovery of myelination.
54. The method of claim 46, wherein rotamase inhibition is achieved using no greater than a 1800 nM concentration of the compound.
55. The method of claim 46, wherein rotamase inhibition is achieved using no greater than a 210 nM concentration of

the compound.

56. The method of claim 46, wherein rotamase inhibition is achieved using no greater than a 103 nM concentration of the compound.

57. The method of claim 46, wherein neurotrophic  $ED_{50}$  is achieved using no greater than a 200 nM concentration of the compound.

58. The method of claim 46, wherein neurotrophic  $ED_{50}$  is achieved using no greater than a 35 nM concentration of the compound.

59. The method of claim 46, wherein neurotrophic  $ED_{50}$  is achieved using no greater than a 1 nM concentration of the compound.

60. A method for preventing neurodegeneration in an animal, comprising:

administering to an animal an effective amount of a non-immunosuppressive pipecolic acid derivative having an affinity for FKBP-type immunophilins to prevent neurodegeneration, wherein the FKBP-type immunophilin exhibits rotamase activity and the non-immunosuppressive pipecolic acid derivative inhibits said rotamase activity of the immunophilin.

61. The method of claim 60, further comprising administering an effective amount of a neurotrophic factor to prevent neurodegeneration selected from the group consisting of neurotrophic growth factor, brain derived growth factor, glial derived growth factor, ciliary neurotrophic factor, and neurotrophin-3.
62. The method of claim 60, wherein the non-immunosuppressive pipecolic acid derivative is Way-124,666.
63. The method of claim 60, wherein the non-immunosuppressive pipecolic acid derivative is Rap-Pa.
64. The method of claim 60, wherein the non-immunosuppressive pipecolic acid derivative is SLB-506.
65. The method of claim 60, wherein the non-immunosuppressive pipecolic acid derivative is selected from the group consisting of compounds 3-84, 86-88 and 90-111.
66. The method of claim 60, wherein the method provides at least 75% regeneration of axon number and at least 35% recovery of myelination.
67. The method of claim 60, wherein the method provides at least 75% regeneration of axon number and at least 50% recovery of myelination.



68. The method of claim 60, wherein rotamase inhibition is achieved using no greater than a 1800 nM concentration of the compound.
69. The method of claim 60, wherein rotamase inhibition is achieved using no greater than a 210 nM concentration of the compound.
70. The method of claim 60, wherein rotamase inhibition is achieved using no greater than a 103 nM concentration of the compound.
71. The method of claim 60, wherein neurotrophic  $ED_{50}$  is achieved using no greater than a 200 nM concentration of the compound.
72. The method of claim 60, wherein neurotrophic  $ED_{50}$  is achieved using no greater than a 35 nM concentration of the compound.
73. The method of claim 60, wherein neurotrophic  $ED_{50}$  is achieved using no greater than a 1 nM concentration of the compound.

1/21




FIG. 1

2/21




FIG. 2A




FIG. 2B

3/21

FIG. 3A





FIG. 3B





FIG. 3C

4/21

**FK-506 AUTORAD**



**FKBP IN SITU**



**CONTROL**

**1 WEEK**

**6 WEEK**

**FIG. 4**

5/21



FIG. 5A




FIG. 5B

FIG. 6

Proximal  Distal





FIG. 8



7/21

FIG. 7A

FKBP IN SITU



Control


FIG. 7C

FK-506 AUTORAD




Control

FKBP IN SITU



Crush

FK-506 AUTORAD



Crush

FIG. 7B

FIG. 7D



8/21

FIG. 9B




FIG. 9D




FIG. 9A





FIG. 9C



9/21

FIG. 10





FIG. 11



10/21




FIG. 12

11/21

FIG. 13A




FIG. 13B



FIG. 13C



FIG. 13D



FIG. 13E



FIG. 13F

12/21

FIG. 14





FIG. 15



14/21




FIG. 16A




FIG. 16B




FIG. 16C




FIG. 16D



16/21




FIG. 16E




FIG. 16F

17/21




FIG. 17A




FIG. 17B






FIG. 17C

18/21




**Example 111 Promotes Neurite Outgrowth in Chick Sensory Neurons**



**1 pM Example 111**






**100 pM Example 111**



**100 nM Example 111**

**FIG. 19**

20/21

**Example 17 Promotes Neurite Outgrowth in Chick Sensory Neuronal Cultures****1 pM Example 17****100 pM Example 17****100 nM Example 17****FIG. 20**

21/21


# Example 102 Elicits Neuronal Outgrowth from Chick Sensory Neurons



**1 pM Example 102**



**100 pM Example 102**



**100 nM Example 102**

**FIG. 21**

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US96/09561

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) :A61K 31/495, 31/50, 31/44, 31/445

US CL :514/248, 293, 300, 318, 326, 330, 302

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/248, 293, 300, 318, 326, 330, 302

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

STN(REGISTRY(STRUCTURE), CA, BIOSIS, MEDLINE, USPATFULL, DRUGU)

search terms: Alzheimer's Disease, Parkinson's Disease, amyotrophic lateral sclerosis, als, neuropathies, brain damage, neurodegeneration, stroke, neural regeneration, nerve growth

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Chemical Abstracts, Volume 120, No. 25 issued 20 June 1994, Lyons et al, "Immunosuppressant FK506 promotes neurite outgrowth in cultures of PC12 cells and sensory ganglia" see page 48, column 1, abstract no. 315423r, Proc. Natl. Acad. Sci. U.S.A., 91(8), 3191-3195, see entire document.	1-56
X	Chemical Abstracts, Volume 121, No. 19 issued 07 November 1994, Sharkey et al, "Immunophilins mediate the neuroprotective effects of FK506 in focal cerebral ischemia" see page 95, column 1, abstract no. 221398e, Nature, 371(6495), 336-339, see entire document.	1-56

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*A* document defining the general state of the art which is not considered to be of particular relevance	X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*E* earlier document published on or after the international filing date	Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	&* document member of the same patent family
*O* document referring to an oral disclosure, use, exhibition or other means	
*P* document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

22 AUGUST 1996

Date of mailing of the international search report

30 SEP 1996

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

WILLIAM JARVIS

Telephone No. (703) 308-1235

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US96/09561

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Chemical Abstracts, Volume 121, No. 19 issued 07 November 1994, Gold et al, "The immunosuppressant FK506 increases functional recovery and nerve regeneration following peripheral nerve injury", see page 105, column 2, abstract no. 221678w, Restor. Neurol. Neurosci., 6(4), 287-296, see entire document.	1-56
X	Chemical Abstracts, Volume 122, No. 10 issued 06 March 1995, Kelly et al, "Macrolide compositions for treatment of amyotrophic lateral sclerosis", see page 659, column 1, abstract no. 114965m, Brit. UK Pat. Appl. GB 2,279,006, pages 1-14, see entire document.	1-56